

Metadata of the article that will be visualized in Online First

Journal Name	Revista Matemática Complutense	
Article Title	Weak optimal controls in coefficients for linear elliptic problems	
Copyright holder	Revista Matemática Complutense This will be the copyright line in the final PDF.	
Corresponding Author	Family name	Buttazzo
	Particle	
	Given Name	Giuseppe
	Suffix	
	Division	Dipartimento di Matematica
	Organization	Università di Pisa
	Address	Largo B. Pontecorvo 5, 56127, Pisa, Italy
	E-mail	buttazzo@dm.unipi.it
Author	Family name	Kogut
	Particle	
	Given Name	Peter
	Given Name	I.
	Suffix	
	Division	Department of Differential Equations
	Organization	Dnipropetrovsk National University
	Address	Naukova str. 13, 49050, Dnipropetrovsk, Ukraine
	E-mail	p.kogut@i.ua
Schedule	Received	7 December 2009
	Revised	
	Accepted	8 February 2010
Abstract	In this paper we study an optimal control problem associated to a linear degenerate elliptic equation with mixed boundary conditions. The equations of this type can exhibit the Lavrentieff phenomenon and non-uniqueness of weak solutions. We adopt the weight function as a control in $L^1(\Omega)$. Using the direct method in the Calculus of variations, we discuss the solvability of this optimal control problem in the class of weak admissible solutions.	
Keywords	Degenerate elliptic equations – Control in coefficients – Weighted Sobolev spaces – Lavrentieff phenomenon – Calculus of variations Mathematics Subject Classification (2000) 35J70 – 49J20 – 49J45 – 93C73	
Footnotes		

Weak optimal controls in coefficients for linear elliptic problems

Giuseppe Buttazzo · Peter I. Kogut

Received: 7 December 2009 / Accepted: 8 February 2010
© Revista Matemática Complutense

Abstract In this paper we study an optimal control problem associated to a linear degenerate elliptic equation with mixed boundary conditions. The equations of this type can exhibit the Lavrentieff phenomenon and non-uniqueness of weak solutions. We adopt the weight function as a control in $L^1(\Omega)$. Using the direct method in the Calculus of variations, we discuss the solvability of this optimal control problem in the class of weak admissible solutions.

Keywords Degenerate elliptic equations · Control in coefficients · Weighted Sobolev spaces · Lavrentieff phenomenon · Calculus of variations

Mathematics Subject Classification (2000) 35J70 · 49J20 · 49J45 · 93C73

1 Introduction

The aim of this work is to study the existence of optimal solutions in coefficients associated to a linear degenerate elliptic equation with mixed boundary condition. By control variable we mean a weight coefficient in the main part of the elliptic operator. The precise answer of existence or none-existence of an L^1 -optimal solutions is heavily depending on the class of admissible controls. Here are the main questions: what is the right setting of optimal control problem with L^1 -controls in coefficients, and what is the right class of admissible solutions to the above problem? Using the

G. Buttazzo (✉)
Dipartimento di Matematica, Università di Pisa, Largo B. Pontecorvo 5, 56127 Pisa, Italy
e-mail: buttazzo@dm.unipi.it

P.I. Kogut
Department of Differential Equations, Dnipropetrovsk National University, Naukova str. 13,
49050 Dnipropetrovsk, Ukraine
e-mail: p.kogut@i.ua

48 direct method in the Calculus of variations, we discuss the solvability of this optimal
49 control problem in the so-called class of weak admissible solutions.

50 In this paper we deal with an optimal control problem in coefficients for boundary
51 value problem of the form

$$\begin{cases} -\operatorname{div}(\rho(x)\nabla y) + y = f & \text{in } \Omega, \\ y = 0 & \text{on } \Gamma_D, \\ \rho(x)\frac{\partial y}{\partial \nu} = 0 & \text{on } \Gamma_N, \end{cases} \quad (1)$$

52
53
54
55
56
57 where $f \in L^2(\Omega)$ is a given function, the boundary of Ω is made of two disjoint
58 parts $\partial\Omega = \Gamma_D \cup \Gamma_N$, and ρ is a measurable non negative weight on a bounded open
59 domain Ω in \mathbb{R}^N .

60 Several physical phenomena related to equilibrium of continuous media are modeled
61 by this elliptic problem. In order to be able to handle with media which possibly
62 are somewhere “perfect” insulators or “perfect” conductors (see [9]) we allow the
63 weight ρ to vanish somewhere or to be unbounded.

64 Though numerous papers (see, for instance, [4, 7, 8, 10, 12, 13, 16] and references
65 therein) are devoted to variational and non variational approaches to problems related
66 to (1), only few papers deal with optimal control problems for degenerate partial dif-
67 ferential equations (see for example [2, 3]). This can be explained by several reasons.
68 Firstly, boundary value problem (1) for locally integrable weight function ρ may ex-
69 hibit the Lavrentieff phenomenon, the non-uniqueness of weak solutions, as well as
70 other surprising consequences. So, in general, the mapping $\rho \mapsto y(\rho)$ can be mul-
71 ti-valued. Besides, the characteristic feature of this problem is the fact that for every
72 admissible control function ρ with properties prescribed above, the weak solutions of
73 (1) belong to the corresponding weighted Sobolev space $W^{1,p}(\Omega, \rho dx)$. In addition,
74 even if the original elliptic equation is non-degenerate, i.e. admissible controls ρ are
75 such that $\rho(x) \geq \alpha > 0$, the majority of optimal control problems in coefficients have
76 no solution (see for instance [11]).

77 The optimal control problem we consider in this paper is closely related to the opti-
78 mal reinforcement of an elastic membrane [3]. Reinforcing an elastic structure sub-
79 jected to a given load is a problem which arises in several applications. The literature
80 on the topic is very wide; for a clear description of the problem from a mechanical
81 point of view and a related bibliography we refer for instance to the beautiful paper
82 by Villaggio [14].

83 In the simplest case when we have an elastic membrane occupying a domain Ω
84 and subjected to a given exterior load $f \in L^2(\Omega)$, the shape u of the membrane in the
85 equilibrium configuration is characterized as the solution of the partial differential
86 equation

$$-\operatorname{div}(\rho(x)\nabla y) + y = f \quad \text{in } \Omega$$

87
88
89 together with the corresponding Dirichlet and Neumann boundary conditions on $\partial\Omega$.
90 The reinforcement of the membrane is usually performed by the addition of suitable
91 stiffeners, whose total amount is prescribed. Mathematically, this is described by a
92 nonnegative coefficient $\rho(x)$ which acts in Ω and is associated with some weight
93 coefficient in the main part of elliptic operator. As a result, the problem of finding
94

an optimal reinforcement for the membrane then consists in the determination of a weight $\rho(x) \geq 0$ which optimizes a given cost functional. In contrast to the paper [3], we do not restrict of our analysis to the particular case of the reinforcement problems. We also do not make use of any relaxations for the original optimal control problem.

2 Notation and preliminaries

Let Ω be a bounded open subset of \mathbb{R}^N ($N \geq 1$) with a Lipschitz boundary. We assume that the boundary of Ω is made of two disjoint parts

$$\partial\Omega = \Gamma_D \cup \Gamma_N$$

with Dirichlet boundary conditions on Γ_D , and Neumann boundary conditions on Γ_N . Let χ_E be the characteristic function of a subset $E \subset \Omega$, i.e. $\chi_E(x) = 1$ if $x \in E$, and $\chi_E(x) = 0$ if $x \notin E$.

Let $C_0^\infty(\mathbb{R}^N; \Gamma_D) = \{\varphi \in C_0^\infty(\mathbb{R}^N) : \varphi = 0 \text{ on } \Gamma_D\}$. The space $W^{1,1}(\Omega; \Gamma_D)$ is the closure of $C_0^\infty(\mathbb{R}^N; \Gamma_D)$ in the classical Sobolev space $W^{1,1}(\Omega)$. For any subset $E \subset \Omega$ we denote by $|E|$ its N -dimensional Lebesgue measure $\mathcal{L}^N(E)$.

Let $\rho : \mathbb{R}^N \rightarrow \mathbb{R}$ be a locally integrable function on \mathbb{R}^N such that $\rho(x) \geq 0$ for a.e. $x \in \mathbb{R}^N$. Then ρ gives rise to a measure on the measurable subsets of \mathbb{R}^N through integration. This measure will also be denoted by ρ . Thus $\rho(E) = \int_E \rho dx$ for measurable sets $E \subset \mathbb{R}^N$.

We will use the standard notation $L^2(\Omega, \rho dx)$ for the set of measurable functions f on Ω such that

$$\|f\|_{L^2(\Omega, \rho dx)} = \left(\int_{\Omega} f^2 \rho dx \right)^{1/2} < +\infty.$$

We say that a locally integrable function $\rho : \mathbb{R}^N \rightarrow \mathbb{R}_+$ is a weight on Ω if

$$\rho + \rho^{-1} \in L^1_{loc}(\mathbb{R}^N). \tag{2}$$

Note that in this case the functions in $L^2(\Omega, \rho dx)$ are Lebesgue integrable on Ω .

To each weight function ρ we may associate two weighted Sobolev spaces:

$$W_\rho = W(\Omega, \rho dx) \quad \text{and} \quad H_\rho = H(\Omega, \rho dx),$$

where W_ρ is the set of functions $y \in W^{1,1}(\Omega; \Gamma_D)$ for which the norm

$$\|y\|_\rho = \left(\int_{\Omega} (y^2 + \rho |\nabla y|^2) dx \right)^{1/2} \tag{3}$$

is finite, and H_ρ is the closure of $C_0^\infty(\Omega; \Gamma_D)$ in W_ρ . Note that due to the estimates

$$\int_{\Omega} |y| dx \leq \left(\int_{\Omega} |y|^2 dx \right)^{1/2} |\Omega|^{1/2} \leq C \|y\|_\rho, \tag{4}$$

$$\int_{\Omega} |\nabla y| dx \leq \left(\int_{\Omega} |\nabla y|^2 \rho dx \right)^{1/2} \left(\int_{\Omega} \rho^{-1} dx \right)^{1/2} \leq C \|y\|_{\rho}, \quad (5)$$

the space W_{ρ} is complete with respect to the norm $\|\cdot\|_{\rho}$. It is clear that $H_{\rho} \subset W_{\rho}$, and W_{ρ}, H_{ρ} are Hilbert spaces. If ρ is bounded between two positive constants, then it is easy to verify that $W_{\rho} = H_{\rho}$. However, for a “typical” weight ρ the space of smooth functions $C_0^{\infty}(\Omega)$ is not dense in W_{ρ} . Hence the identity $W_{\rho} = H_{\rho}$ is not always valid (for the corresponding examples we refer to [5, 15]).

Weak compactness criterion in $L^1(\Omega)$ Throughout the paper we will often use the concepts of the weak and strong convergence in $L^1(\Omega)$. Let $\{a_{\varepsilon}\}_{\varepsilon>0}$ be a sequence in $L^1(\Omega)$. We recall that $\{a_{\varepsilon}\}_{\varepsilon>0}$ is called equi-integrable if for any $\delta > 0$ there is $\tau = \tau(\delta)$ such that $\int_S |a_{\varepsilon}| dx < \delta$ for every measurable subset $S \subset \Omega$ of Lebesgue measure $|S| < \tau$. Then the following assertions are equivalent:

- (i) a sequence $\{a_{\varepsilon}\}_{\varepsilon>0}$ is weakly compact in $L^1(\Omega)$;
- (ii) the sequence $\{a_{\varepsilon}\}_{\varepsilon>0}$ is equi-integrable;
- (iii) given $\delta > 0$ there exists $\lambda = \lambda(\delta)$ such that $\sup_{\varepsilon>0} \int_{\{|a_{\varepsilon}|>\lambda\}} |a_{\varepsilon}| dx < \delta$.

Theorem 2.1 (Lebesgue’s theorem) *If a sequence $\{a_{\varepsilon}\}_{\varepsilon>0} \subset L^1(\Omega)$ is equi-integrable and $a_{\varepsilon} \rightarrow a$ almost everywhere in Ω then $a_{\varepsilon} \rightarrow a$ in $L^1(\Omega)$.*

Radon measures By a nonnegative Radon measure on Ω we mean a nonnegative Borel measure which is finite on every compact subset of Ω . The space of all nonnegative Radon measures on Ω will be denoted by $M_+(\Omega)$. According to the Riesz theory, each Radon measure $\mu \in M_+(\Omega)$ can be interpreted as element of the dual of the space $C_0(\Omega)$ of all continuous functions vanishing at infinity. Let $M(\Omega; \mathbb{R}^N)$ denotes the space of all \mathbb{R}^N -valued Borel measures. Then $\mu = (\mu_1, \dots, \mu_N) \in M(\Omega; \mathbb{R}^N) \Leftrightarrow \mu_i \in C_0'(\Omega), i = 1, \dots, N$.

If μ is a nonnegative Radon measure on Ω , we will use $L^r(\Omega, d\mu), 1 \leq r \leq \infty$, to denote the usual Lebesgue space with respect to the measure μ with the corresponding norm $\|f\|_{L^r(\Omega, d\mu)} = \left(\int_{\Omega} |f(x)|^r d\mu \right)^{1/r}$.

Functions with bounded variation Let $f : \Omega \rightarrow \mathbb{R}$ be a function of $L^1(\Omega)$. Define

$$\int_{\Omega} |Df| = \sup \left\{ \int_{\Omega} f \operatorname{div} \varphi dx : \varphi = (\varphi_1, \dots, \varphi_N) \in C_0^1(\Omega; \mathbb{R}^N), |\varphi(x)| \leq 1 \text{ for } x \in \Omega \right\},$$

where $\operatorname{div} \varphi = \sum_{i=1}^N \frac{\partial \varphi_i}{\partial x_i}$. According to the Radon-Nikodym theorem, if $\int_{\Omega} |Df| < +\infty$ then the distribution Df is a measure and there exist a vector-valued function $\nabla f \in [L^1(\Omega)]^N$ and a measure $D_s f$, singular with respect to the N -dimensional Lebesgue measure $\mathcal{L}^N \lfloor \Omega$ restricted to Ω , such that

$$Df = \nabla f \mathcal{L}^N \lfloor \Omega + D_s f.$$

Definition 2.2 A function $f \in L^1(\Omega)$ is said to have a bounded variation in Ω if $\int_{\Omega} |Df| < +\infty$. By $BV(\Omega)$ we denote the space of all functions in $L^1(\Omega)$ with bounded variation.

Under the norm

$$\|f\|_{BV(\Omega)} = \|f\|_{L^1(\Omega)} + \int_{\Omega} |Df|,$$

$BV(\Omega)$ is a Banach space. It is well-known the following compactness result for BV -functions:

Proposition 2.3 *The uniformly bounded sets in BV -norm are relatively compact in $L^1(\Omega)$.*

Definition 2.4 A sequence $\{f_k\}_{k=1}^{\infty} \subset BV(\Omega)$ weakly converges to some $f \in BV(\Omega)$, and we write $f_k \rightharpoonup f$ iff the two following conditions hold: $f_k \rightarrow f$ strongly in $L^1(\Omega)$, and $Df_k \rightharpoonup Df$ weakly* in $M(\Omega; \mathbb{R}^N)$.

In the proposition below we give a compactness result related to this convergence, together with the lower semicontinuity property (see [6]):

Proposition 2.5 *Let $\{f_k\}_{k=1}^{\infty}$ be a sequence in $BV(\Omega)$ strongly converging to some f in $L^1(\Omega)$ and satisfying $\sup_{k \in \mathbb{N}} \int_{\Omega} |Df_k| < +\infty$. Then*

- (i) $f \in BV(\Omega)$ and $\int_{\Omega} |Df| \leq \liminf_{k \rightarrow \infty} \int_{\Omega} |Df_k|$;
- (ii) $f_k \rightharpoonup f$ in $BV(\Omega)$.

Convergence in variable spaces Let $\{\mu_k\}_{k \in \mathbb{N}}$, μ be Radon measures such that $\mu_k \xrightarrow{*} \mu$ in $M_+(\Omega)$, i.e.,

$$\lim_{k \rightarrow \infty} \int_{\Omega} \varphi d\mu_k = \int_{\Omega} \varphi d\mu \quad \forall \varphi \in C_0(\mathbb{R}^N), \quad (6)$$

where $C_0(\mathbb{R}^N)$ is the space of all compactly supported continuous functions. The typical example of such measures is

$$d\mu_k = \rho_k(x) dx, \quad d\mu = \rho(x) dx, \quad \text{where } 0 \leq \rho_k \rightharpoonup \rho \text{ in } L^1(\Omega).$$

Let us recall the definition and main properties of convergence in the variable L^2 -space (see [15]).

1. A sequence $\{v_k \in L^2(\Omega, d\mu_k)\}$ is called bounded if

$$\limsup_{k \rightarrow \infty} \int_{\Omega} |v_k|^2 d\mu_k < +\infty.$$

2. A bounded sequence $\{v_k \in L^2(\Omega, d\mu_k)\}$ converges weakly to $v \in L^2(\Omega, d\mu)$ if

$$\lim_{k \rightarrow \infty} \int_{\Omega} v_k \varphi d\mu_k = \int_{\Omega} v \varphi d\mu \quad \forall \varphi \in C_0^{\infty}(\Omega),$$

and it is written as $v_k \rightharpoonup v$ in $L^2(\Omega, d\mu_k)$.

3. The strong convergence $v_k \rightarrow v$ in $L^2(\Omega, d\mu_k)$ means that $v \in L^2(\Omega, d\mu)$ and

$$\lim_{k \rightarrow \infty} \int_{\Omega} v_k z_k d\mu_k = \int_{\Omega} v z d\mu \quad \text{as } z_k \rightarrow z \text{ in } L^2(\Omega, d\mu_k). \quad (7)$$

The following convergence properties in variable spaces hold:

(a) *Compactness*: if a sequence is bounded in $L^2(\Omega, d\mu_k)$, then this sequence is compact in the sense of the weak convergence;

(b) *Lower semicontinuity*: if $v_k \rightharpoonup v$ in $L^2(\Omega, d\mu_k)$, then

$$\liminf_{\varepsilon \rightarrow 0} \int_{\Omega} |v_k|^2 d\mu_k \geq \int_{\Omega} v^2 d\mu; \quad (8)$$

(c) *Strong convergence*: $v_k \rightarrow v$ if and only if $v_k \rightharpoonup v$ in $L^2(\Omega, d\mu_k)$ and

$$\lim_{k \rightarrow \infty} \int_{\Omega} |v_k|^2 d\mu_k = \int_{\Omega} v^2 d\mu. \quad (9)$$

3 Setting of the optimal control problem

Let $m \in \mathbb{R}_+$ be some positive value, and let ξ_1, ξ_2 be given elements of $L^1(\Omega)$ satisfying the conditions

$$\xi_1(x) \leq \xi_2(x) \quad \text{a.e. in } \Omega, \quad \xi_1^{-1} \in L^1(\Omega). \quad (10)$$

To introduce the class of admissible BV -controls we adopt the following concept:

Definition 3.1 We say that a nonnegative weight ρ is an admissible control to the boundary value problem

$$-\operatorname{div}(\rho(x)\nabla y) + y = f \quad \text{in } \Omega, \quad (11)$$

$$y = 0 \quad \text{on } \Gamma_D, \quad \rho(x) \frac{\partial y}{\partial \nu} = 0 \quad \text{on } \Gamma_N, \quad (12)$$

(it is written as $\rho \in \mathcal{R}_{ad}$) if

$$\rho \in BV(\Omega), \quad \int_{\Omega} \rho dx = m, \quad \xi_1(x) \leq \rho(x) \leq \xi_2(x) \quad \text{a.e. in } \Omega. \quad (13)$$

Here $f \in L^2(\Omega)$ is a given function.

Hereinafter we assume that the set \mathcal{R}_{ad} is nonempty.

Remark 3.2 In view of the property (10), we have the boundary value problem for the degenerate elliptic equation. It means that for some admissible controls $\rho \in \mathcal{R}_{ad}$ the boundary value problem (11)–(12) can exhibit the Lavrentieff phenomenon, the nonuniqueness of the weak solutions as well as other surprising consequences.

The optimal control problem we consider in this paper is to minimize the discrepancy between a given distribution $y_d \in L^2(\Omega)$ and the solution of boundary valued problem (11)–(12) by choosing an appropriate weight function $\rho \in \mathcal{R}_{ad}$. More precisely, we are concerned with the following optimal control problem

$$\text{Minimize } \left\{ I(\rho, y) = \int_{\Omega} |y(x) - y_d(x)|^2 dx + \int_{\Omega} |\nabla y(x)|_{\mathbb{R}^N}^2 \rho dx + \int_{\Omega} |D\rho| \right\} \quad (14)$$

subject to the constraints (11)–(13).

Definition 3.3 We say that a function $y = y(\rho, f) \in W_{\rho}$ is a weak solution to the boundary value problem (11)–(12) for a fixed control $\rho \in \mathcal{R}_{ad}$ if the integral identity

$$\int_{\Omega} (\rho \nabla y \cdot \nabla \varphi + y\varphi) dx = \int_{\Omega} f\varphi dx \quad (15)$$

holds for any $\varphi \in C_0^{\infty}(\Omega; \Gamma_D)$.

It is clear that the question of uniqueness of a weak solution leads us to the problem of density of the subspace of smooth functions $C_0^{\infty}(\Omega; \Gamma_D)$ in W_{ρ} . However, as was indicated in [16], for a “typical” weight function ρ the subspace $C_0^{\infty}(\Omega; \Gamma_D)$ is not dense in W_{ρ} , and hence there is no uniqueness of weak solutions (for more details and other types of solutions we refer to [1, 10, 15, 16]). Thus the mapping $\rho \mapsto y(\rho, f)$ is multivalued, in general. Taking this fact into account, we introduce the set

$$\Xi_W = \{(\rho, y) \mid \rho \in \mathcal{R}_{ad}, y \in W_{\rho}, (\rho, y) \text{ are related by (15)}\}. \quad (16)$$

Note that the set Ξ_W is always nonempty. Indeed, let V_{ρ} be some intermediate space with $H_{\rho} \subset V_{\rho} \subset W_{\rho}$. We say that a function $y = y(\rho, f) \in V_{\rho}$ is a V_{ρ} -solution or variational solution to the boundary value problem (11)–(12) if the integral identity (15) holds for every test function $\varphi \in V_{\rho}$. Hence, in this case the energy equality

$$\int_{\Omega} (|\nabla y|^2 \rho + y^2) dx = \int_{\Omega} f y dx \quad (17)$$

must be valid. Since

$$\left| \int_{\Omega} f y dx \right| \leq \left(\int_{\Omega} f^2 dx \right)^{1/2} \left(\int_{\Omega} y^2 dx \right)^{1/2} \leq C \|y\|_{\rho}$$

for every fixed $f \in L^2(\Omega)$, it follows that the existence and uniqueness of a V_{ρ} -solution are the direct consequence of the Riesz Representation Theorem. Thus every variational solution is also a weak solution to the problem (11)–(12). Hence $\Xi_W \neq \emptyset$ and therefore the corresponding minimization problem

$$\inf_{(\rho, y) \in \Xi_W} I(\rho, y) \quad (18)$$

is regular. In view of this, we adopt the following concept.

Definition 3.4 We say that a pair $(\rho^0, y^0) \in L^1(\Omega) \times W^{1,1}(\Omega; \Gamma_D)$ is a weak optimal solution to the problem (13)–(14) if (ρ^0, y^0) is a minimizer for (18).

4 Existence of weak optimal solutions

Our prime interest in this section deals with the solvability of optimal control problem (13)–(14) in the class of weak solutions. To begin with, we make use of the following results. Let $\{(\rho_k, y_k) \in \Xi_W\}_{k \in \mathbb{N}}$ be any sequence of weak admissible solutions.

Lemma 4.1 Let $\{\rho_k\}_{k \in \mathbb{N}}$ be a sequence in \mathcal{R}_{ad} such that $\rho_k \rightarrow \rho$ in $L^1(\Omega)$. Then

$$(\rho_k)^{-1} \rightarrow \rho^{-1} \quad \text{in the variable space } L^2(\Omega, \rho_k dx).$$

Proof To prove this result we use some ideas of [16]. By the properties of the set of admissible controls \mathcal{R}_{ad} , we have $\rho_k^{-1} \leq \xi_1^{-1}$ for every $k \in \mathbb{N}$, hence the sequence $\{\rho_k^{-1}\}_{k \in \mathbb{N}}$ is equi-integrable on Ω . Note that, up to a subsequence, we have $\rho_k \rightarrow \rho$ a.e. in Ω . Since $\xi_2^{-1} \leq \rho_k^{-1} \leq \xi_1^{-1}$, Lebesgue Theorem implies

$$\rho_k^{-1} \rightarrow \rho^{-1} \quad \text{in } L^1(\Omega). \tag{19}$$

Let $\varphi \in C_0^\infty(\Omega)$ be a fixed function. Then the equality

$$\int_{\Omega} \rho_k^{-1} \varphi \rho_k dx = \int_{\Omega} \varphi dx = \int_{\Omega} \rho^{-1} \varphi \rho dx \quad \forall k \in \mathbb{N}$$

leads us to the weak convergence $\rho_k^{-1} \rightharpoonup \rho^{-1}$ in $L^2(\Omega, \rho_k dx)$. Taking into account the strong convergence $\rho_k^{-1} \rightarrow \rho^{-1}$ in $L^1(\Omega)$ and the fact that Ω is a bounded domain, we get

$$\lim_{k \rightarrow \infty} \int_{\Omega} |\rho_k|^{-2} \rho_k dx = \lim_{k \rightarrow \infty} \int_{\Omega} \rho_k^{-1} dx = \int_{\Omega} \rho^{-1} dx = \int_{\Omega} |\rho|^{-2} \rho dx.$$

Hence, by the strong convergence criterium in the variable space $L^2(\Omega, \rho_k dx)$, we come to the required conclusion. \square

Our next step deals with the study of topological properties of the set of weak admissible solutions Ξ_W to the problem (11)–(14).

Definition 4.2 A sequence $\{(\rho_k, y_k) \in \Xi_W\}_{k \in \mathbb{N}}$ is called bounded if

$$\sup_{k \in \mathbb{N}} [\|\rho_k\|_{BV(\Omega)} + \|y_k\|_{L^2(\Omega)} + \|\nabla y_k\|_{L^2(\Omega, \rho_k dx)^N}] < +\infty.$$

Definition 4.3 We say that a bounded sequence $\{(\rho_k, y_k) \in \Xi_W\}_{k \in \mathbb{N}}$ of the weak admissible solutions τ -converges to a pair $(\rho, y) \in BV(\Omega) \times W^{1,1}(\Omega)$ if

- (a) $\rho_k \rightarrow \rho$ in $BV(\Omega)$;

- (d) $y_k \rightharpoonup y$ weakly in $L^2(\Omega)$;
 (e) $\nabla y_k \rightharpoonup \nabla y \ni L^2(\Omega, \rho dx)^N$ in the variable space $L^2(\Omega, \rho_k dx)^N$.

Note that due to assumptions (10), (13), and estimates like (4)–(5), the inclusion $y \in W^{1,1}(\Omega)$ is obvious.

Lemma 4.4 *Let $\{(\rho_k, y_k) \in \Xi_W\}_{k \in \mathbb{N}}$ be a bounded sequence. Then there is a pair $(\rho, y) \in BV(\Omega) \times W^{1,1}(\Omega)$ such that, up to a subsequence, $(\rho_k, y_k) \xrightarrow{\tau} (\rho, y)$ and $y \in W_\rho$.*

Proof By Proposition 2.3 and the compactness criterium of the weak convergence in variable spaces, there exist a subsequence of $\{(\rho_k, y_k) \in \Xi_W\}_{k \in \mathbb{N}}$, still denoted by the same indices, and functions $\rho \in BV(\Omega)$, $y \in L^2(\Omega)$, and $v \in L^2(\Omega, \rho dx)^N$ such that

$$\rho_k \rightarrow \rho \quad \text{in } L^1(\Omega), \tag{20}$$

$$y_k \rightharpoonup y \quad \text{in } L^2(\Omega), \quad \nabla y_k \rightharpoonup v \quad \text{in the variable space } L^2(\Omega, \rho_k dx). \tag{21}$$

Let us show that $y \in W^{1,1}(\Omega)$, and $v = \nabla y$. Since $\xi_1 \leq \rho_k \leq \xi_2$ for every $k \in \mathbb{N}$, (20) and Lemma 4.1 imply the property (see (19))

$$\rho_k^{-1} \rightarrow \rho^{-1} \quad \text{in } L^1(\Omega), \quad \xi_1 \leq \rho \leq \xi_2 \quad \text{a.e. in } \Omega. \tag{22}$$

This yields that the sequence $\{\nabla y_k\}_{k \in \mathbb{N}}$ is weakly compact in $L^1(\Omega)^N$. Indeed, the property of its equi-integrability immediately follows from the inequality

$$\int_E |\nabla y_k| dx \leq \left(\int_E \rho_k^{-1} dx \right)^{1/2} \left(\int_\Omega |\nabla y_k|^2 \rho_k dx \right)^{1/2} \leq C \left(\int_E \rho_k^{-1} dx \right)^{1/2}.$$

As a result, using the strong convergence $(\rho_k)^{-1} \rightarrow \rho^{-1}$ in the variable space $L^2(\Omega, \rho_k dx)$ (see Lemma 4.1) and its properties, we obtain

$$\begin{aligned} \lim_{k \rightarrow \infty} \int_\Omega \nabla y_k \cdot \psi dx &= \lim_{k \rightarrow \infty} \int_\Omega \rho_k^{-1} \nabla y_k \cdot \psi \rho_k dx \\ &= \int_\Omega \rho^{-1} v \cdot \psi \rho dx = \int_\Omega v \cdot \psi dx \end{aligned}$$

for all $\psi \in C_0^\infty(\Omega)^N$. Thus $\nabla y_k \rightharpoonup v$ in $L^1(\Omega)^N$. This implies that $y \in W^{1,1}(\Omega)$ and $\nabla y = v$. The inclusion $y \in W_\rho$ immediately follows from (20)–(21). \square

Theorem 4.5 *For every $f \in L_{loc}^2(\mathbb{R}^N)$ the set Ξ_W is sequentially closed with respect to the τ -convergence.*

Proof Let $\{(\rho_k, y_k)\}_{k \in \mathbb{N}} \subset \Xi_W$ be a bounded τ -convergent sequence of weak admissible pairs to the optimal control problem (11)–(14). Let (ρ_0, y_0) be its τ -limit. Our aim is to prove that $(\tau_0, y_0) \in \Xi_W$. By Lemma 4.4 we have

$$\rho_k \rightarrow \rho_0 \quad \text{in } L^1(\Omega), \quad \rho_0 \in BV(\Omega), \quad \xi_1 \leq \rho_0 \leq \xi_2 \quad \text{a.e. in } \Omega. \tag{23}$$

Then passing to the limit as $k \rightarrow \infty$ in the relation $\int_{\Omega} \rho_k dx = m$, we conclude that $\rho_0 \in \mathcal{R}_{ad}$, i.e. the limit weight function ρ_0 is an admissible control.

It remains to show that the pair (ρ_0, y_0) is related by the integral identity (15) for all $\varphi \in C_0^\infty(\Omega; \Gamma_D)$. For every fixed $k \in \mathbb{N}$ we denote by $(\widehat{\rho}_k, \widehat{y}_k) \in BV_{loc}(\mathbb{R}^N) \times W_{loc}^{1,1}(\mathbb{R}^N)$ an extension of the functions $(\rho_k, y_k) \in \Xi_W$ to the whole of space \mathbb{R}^N such that the sequence $\{(\widehat{\rho}_k, \widehat{y}_k)\}_{k \in \mathbb{N}}$ satisfies the properties:

$$\widehat{\rho}_k \in BV(Q), \quad \xi_1 \leq \widehat{\rho}_k \leq \xi_2 \quad \text{a.e. in } Q, \quad (24)$$

$$\sup_{k \in \mathbb{N}} [\|\widehat{\rho}_k\|_{BV(Q)} + \|\widehat{y}_k\|_{L^2(Q)} + \|\nabla \widehat{y}_k\|_{L^2(Q, \widehat{\rho}_k dx)^N}] < +\infty \quad (25)$$

for any bounded domain Q in \mathbb{R}^N . Hence, as done in Lemma 4.4 it can be proved that for every bounded domain $Q \subset \mathbb{R}^N$ there exist functions $\widehat{\rho}_0 \in BV(Q)$ and $\widehat{y}_0 \in W_{\widehat{\rho}_0}$ such that

$$\widehat{\rho}_k \rightarrow \widehat{\rho}_0 \quad \text{in } L^1(Q), \quad \widehat{y}_k \rightarrow \widehat{y}_0 \quad \text{in } L^2(Q), \quad (26)$$

$$\nabla \widehat{y}_k \rightharpoonup \nabla \widehat{y}_0 \ni L^2(\Omega, \widehat{\rho}_0 dx)^N \quad \text{in the variable space } L^2(\Omega, \widehat{\rho}_k dx)^N. \quad (27)$$

It is important to note that in this case we have

$$\widehat{y}_0 = y_0 \quad \text{and} \quad \widehat{\rho}_0 = \rho_0 \quad \text{a.e. in } \Omega. \quad (28)$$

In what follows, we rewrite the integral identity (15) in the equivalent form

$$\int_{\mathbb{R}^N} (\nabla \widehat{y}_k \cdot \nabla \varphi \widehat{\rho}_k + \widehat{y}_k \varphi) \chi_{\Omega}(x) dx = \int_{\mathbb{R}^N} f \varphi \chi_{\Omega}(x) dx \quad \forall \varphi \in C_0^\infty(\mathbb{R}^N, \Gamma_D), \quad (29)$$

and pass to the limit in (29) as $k \rightarrow \infty$. Using the properties (26)–(27), and the fact that $\chi_{\Omega} \rightarrow \chi_{\Omega}$ strongly in the variable space $L^2(Q, \widehat{\rho}_k dx)$, i.e.

$$\int_{\mathbb{R}^N} \chi_{\Omega}^2 \widehat{\rho}_k dx = \int_{\mathbb{R}^N} \chi_{\Omega} \widehat{\rho}_k dx \rightarrow \int_{\mathbb{R}^N} \chi_{\Omega} \widehat{\rho}_0 dx = \int_{\mathbb{R}^N} \chi_{\Omega}^2 \widehat{\rho}_0 dx$$

we obtain

$$\int_{\mathbb{R}^N} (\nabla \widehat{y}_0 \cdot \nabla \varphi \widehat{\rho}_0 + \widehat{y}_0 \varphi) \chi_{\Omega}(x) dx = \int_{\mathbb{R}^N} f \varphi \chi_{\Omega}(x) dx \quad \forall \varphi \in C_0^\infty(\mathbb{R}^N, \Gamma_D)$$

which is equivalent to

$$\int_{\Omega} (\nabla \widehat{y}_0 \cdot \nabla \varphi \widehat{\rho}_0 + \widehat{y}_0 \varphi) dx = \int_{\Omega} f \varphi dx \quad \forall \varphi \in C_0^\infty(\Omega, \Gamma_D).$$

As a result, taking into account (28) and the fact that $\widehat{y}_0 \in W_{\widehat{\rho}_0}$ (by Lemma 4.4), we conclude: y_0 is a weak solution to the boundary valued problem (11)–(12) under $\rho = \rho_0$. Thus the τ -limit pair (τ_0, y_0) belongs to set Ξ_W , and this concludes the proof. \square

We are now in a position to state the existence of weak optimal pairs to the problem (11)–(14).

Theorem 4.6 Let $\xi_1 \in L^1_{loc}(\mathbb{R}^N)$ and $\xi_2 \in L^1_{loc}(\mathbb{R}^N)$ be such that $\xi_1 \leq \xi_2$ a.e. in \mathbb{R}^N and $\xi_1^{-1} \in L^1_{loc}(\mathbb{R}^N)$. Let $f \in L^2_{loc}(\mathbb{R}^N)$ and $y_d \in L^2(\Omega)$ be given functions, and assume that $\mathcal{R}_{ad} \neq \emptyset$. Then the optimal control problem (11)–(14) admits at least one weak solution

$$(\rho^{opt}, y^{opt}) \in \Xi_W \subset L^1(\Omega) \times W^{1,1}(\Omega, \Gamma_D), \quad y^{opt} \in W_{\rho^{opt}}.$$

Proof Since the set of admissible controls \mathcal{R}_{ad} is nonempty the minimization problem (18) is regular (i.e. $\Xi_W \neq \emptyset$). Let $\{(\rho_k, y_k) \in \Xi_W\}_{k \in \mathbb{N}}$ be a minimizing sequence to (18). From the inequality

$$\inf_{(\rho, y) \in \Xi_W} I(\rho, y) = \lim_{k \rightarrow \infty} \left[\int_{\Omega} |y_k(x) - y_d(x)|^2 dx + \int_{\Omega} |\nabla y_k(x)|^2 \rho_k dx + \int_{\Omega} |D\rho_k| \right] < +\infty, \quad (30)$$

there is a constant $C > 0$ such that

$$\sup_{k \in \mathbb{N}} \|y_k\|_{L^2(\Omega)} \leq C, \quad \sup_{k \in \mathbb{N}} \|\nabla y_k\|_{L^2(\Omega, \rho_k dx)^N} \leq C, \quad \int_{\Omega} |D\rho_k| \leq C.$$

Hence, in view of the definition of the class of admissible controls \mathcal{R}_{ad} , the sequence $\{(\rho_k, y_k) \in \Xi_W\}_{k \in \mathbb{N}}$ is bounded in the sense of Definition 4.2. Hence, by Lemma 4.4 there exist functions $\rho^* \in BV(\Omega)$ and $y^* \in W_{\rho^*}$ such that, up to a subsequence, $(\rho_k, y_k) \xrightarrow{\tau} (\rho^*, y^*)$. Since the set Ξ_W is sequentially closed with respect to the τ -convergence (see Theorem 4.5), it follows that the τ -limit pair (ρ^*, y^*) is an admissible weak solution to the optimal control problem (11)–(14) (i.e. $(\rho^*, y^*) \in \Xi_W$). To conclude the proof it is enough to observe that by property (8) and Proposition 2.5, the cost functional I is sequentially lower τ -semicontinuous. Thus

$$I(\rho^*, y^*) \leq \liminf_{k \rightarrow \infty} I(\rho_k, y_k) = \inf_{(\rho, y) \in \Xi_W} I(\rho, y).$$

Hence (ρ^*, y^*) is a weak optimal pair, and we come to the required conclusion. \square

References

1. Boccardo, L., Gallouët, T., Marcellini, P.: Anisotropic equations in L^1 . *Differ. Integral Equ.* **9**, 209–212 (1996)
2. Bouchitte, G., Buttazzo, G.: Characterization of optimal shapes and masses through Monge-Kantorovich equation. *J. Eur. Math. Soc.* **3**, 139–168 (2001)
3. Buttazzo, G., Varchon, N.: On the optimal reinforcement of an elastic membrane. *Riv. Mat. Univ. Parma* **4**(7), 115–125 (2005)
4. Caldiroli, P., Musina, R.: On a variational degenerate elliptic problem. *Nonlinear Differ. Equ. Appl.* **7**, 187–199 (2000)
5. Chiadò Piat, V., Serra Cassano, F.: Some remarks about the density of smooth functions in weighted Sobolev spaces. *J. Convex Anal.* **1**(2), 135–142 (1994)
6. Giusti, E.: *Minimal Surfaces and Functions of Bounded Variation*. Birkhäuser, Boston (1984)

- 518 7. Chabrowski, J.: Degenerate elliptic equation involving a subcritical Sobolev exponent. *Port. Math.* **53**,
519 167–177 (1996)
- 520 8. Cirmi, G.R., Porzio, M.M.: L^∞ -solutions for some nonlinear degenerate elliptic and parabolic equa-
521 tions. *Ann. Mat. Pura Appl.* **169**, 67–86 (1995)
- 522 9. Dautray, R., Lions, J.L.: *Mathematical Analysis and Numerical Methods for Science and Technology. Physical Origins and Classical Methods*, vol. 1. Springer, Berlin (1985)
- 523 10. Kovalevsky, A.A., Gorban, Yu.S.: Degenerate anisotropic variational inequalities with L^1 -data.
524 C.R.A.S. Paris, Sér. I **345**, 441–444 (2007)
- 525 11. Murat, F.: Un contre-exemple pour le problème de contrôle dans les coefficients. C.R.A.S. Paris, Sér.
526 A **273**, 708–711 (1971)
- 527 12. Murthy, M.K.V., Stampacchia, V.: Boundary problems for some degenerate elliptic operators. *Ann.*
528 *Mat. Pura Appl.* **5**(4), 1–122 (1968)
- 529 13. Pastukhova, S.E.: Degenerate equations of monotone type: Lavrentev phenomenon and attainability
530 problems. *Sb. Math.* **198**(10), 1465–1494 (2007)
- 531 14. Villaggio, P.: Calcolo delle variazioni e teoria delle strutture. *Boll. Univ. Mat. Ital. Ser. VIII VII-A*,
532 49–76 (2004)
- 533 15. Zhikov, V.V.: Weighted Sobolev spaces. *Sb. Math.* **189**(8), 27–58 (1998)
- 534 16. Zhikov, V.V., Pastukhova, S.E.: Homogenization of degenerate elliptic equations. *Sib. Math. J.* **49**(1),
535 80–101 (2006)